

ON THE GEOMETRY OF CLOSED G_2 -STRUCTURES

RICHARD CLEYTON AND STEFAN IVANOV

ABSTRACT. We give an answer to a question posed recently by R.Bryant, namely we show that a compact 7-dimensional manifold equipped with a G_2 -structure with closed fundamental form is Einstein if and only if the Riemannian holonomy of the induced metric is contained in G_2 . This could be considered to be a G_2 analogue of the Goldberg conjecture in almost Kähler geometry. The result was generalized by R.L.Bryant to closed G_2 -structures with too tightly pinched Ricci tensor. We extend it in another direction proving that a compact G_2 -manifold with closed fundamental form and divergence-free Weyl tensor is a G_2 -manifold with parallel fundamental form. We introduce a second symmetric Ricci-type tensor and show that Einstein conditions applied to the two Ricci tensors on a closed G_2 -structure again imply that the induced metric has holonomy group contained in G_2 .

CONTENTS

1. Introduction	1
2. General properties of G_2 -structures	3
3. Ricci tensors on G_2 -manifold	4
4. Closed G_2 -structures	5
5. Curvature of closed G_2 -structures	6
6. An integral formula on closed G_2 manifold	9
7. Proof of the Main Theorem	12
References	13

1. INTRODUCTION

A 7-dimensional Riemannian manifold is called a G_2 -manifold if its structure group reduces to the exceptional Lie group G_2 . The existence of a G_2 -structure is equivalent to the existence of a non-degenerate three-form on the manifold, sometimes called the fundamental form of the G_2 -manifold. From the purely topological point of view, a 7-dimensional paracompact manifold is a G_2 -manifold if and only if it is an oriented spin manifold [23].

In [11], Fernandez and Gray divide G_2 -manifolds into 16 classes according to how the covariant derivative of the fundamental three-form behaves with respect to its decomposition into G_2 irreducible components (see also [7]). If the fundamental form is parallel with respect to the Levi-Civita connection then the Riemannian holonomy group is contained in G_2 , we will say that the G_2 -manifold or the G_2 -structure on the manifold is *parallel*. In this case the induced metric on the G_2 -manifold is Ricci-flat, a fact first observed by Bonan [1]. It was shown by Gray [17] (see also [2, 25]) that a G_2 -manifold is parallel precisely when the fundamental form is harmonic. The first examples of complete parallel

G_2 -manifolds were constructed by Bryant and Salamon [5, 16]. Compact examples of parallel G_2 -manifolds were obtained first by Joyce [19, 20, 21] and recently by Kovalev [22]. Compact parallel G_2 -manifold will be referred to as *Joyce spaces*. Examples of G_2 -manifolds in other Fernandez-Gray classes may be found in [10, 6]. A central point in our argument is that the Riemannian scalar curvature of a G_2 -manifold may be expressed in terms of the fundamental form and its derivatives and furthermore the scalar curvature carries a definite sign for certain classes of G_2 -manifolds [13, 3].

The geometry of G_2 -structures has also attracted much attention from physicists. The central issue in physics is that connections with holonomy contained in G_2 plays a rôle in string theory [14, 9, 24, 15, 18]. The G_2 -connections admitting three-form torsion have been of particular interest.

In the present paper we are interested in the geometry of closed G_2 -structures i.e., G_2 -manifolds with closed fundamental form (sometimes these spaces are called almost G_2 -manifolds or calibrated G_2 -manifolds). In the sense of the Fernandez-Gray classes, this is complementary to the physicists requirement of three-form torsion [12]. Compact examples of closed G_2 -manifolds were presented by Fernandez [10]. Supersymmetric string solutions on closed G_2 -manifolds were investigated in [9], topological quantum field theory on closed G_2 -manifolds were discussed in [24]. Robert Bryant shows in [3] that if the scalar curvature of a closed G_2 -structure is non-negative then the G_2 -manifold is parallel. The question whether there are closed G_2 -structure which are Einstein but not Ricci-flat then naturally arises. We investigate this question of Bryant in the compact and in the non-compact cases.

In the first version of the present article [8] we answered negatively to the Bryant's question, namely, we proved that there are no closed Einstein G_2 - structures (other than the parallel ones) on compact 7-manifold. In [4] R.L.Bryant generalized this non-existence result for closed G_2 -structures on compact 7-manifold whose Ricci tensor is too tightly pinched.

In the present article we obtain non-existence result involving third derivatives of the fundamental form. Namely, we prove the following

Main Theorem. *A compact G_2 -manifold with closed fundamental form and harmonic Weyl tensor (divergence-free Weyl tensor) is a Joyce space.*

The second Bianchi identity leads to

Corollary 1.1. *A compact G_2 -manifold with closed fundamental form and harmonic curvature (divergence-free curvature tensor) is a Joyce space.*

Corollary 1.2. *A compact Einstein G_2 -manifold with closed fundamental form is a Joyce space.*

The latter may be considered to be a G_2 analogue of the Goldberg conjecture in almost Kähler geometry (see e.g. [9]).

The representation theory of G_2 gives rise to a second symmetric Ricci type tensor on G_2 -manifolds. Therefore one may consider two complementary Einstein equations. We find a connection between the two Ricci tensors and show in Theorem 5.7, with no compactness assumption, that if both Einstein conditions hold simultaneously on a G_2 -manifold with closed fundamental form then the fundamental form is parallel.

Our main tool is the canonical connection of a G_2 -structure and its curvature. We will show that the Ricci tensor of the canonical connection is proportional to the Riemannian Ricci tensor. This leads to the corollary that a compact G_2 -manifold with closed fundamental form which is Einstein with respect to the canonical connection is a Joyce space.

Our main technical tool is an integral formula which holds on any compact G_2 -manifold with closed fundamental form. We derive the Main Theorem as a consequence of a more general result, Theorem 7.1, which shows that the vanishing of the Λ_7^2 -part of the divergence of the Weyl tensor implies that a closed G_2 -structure is parallel on a compact 7-manifold.

Acknowledgements. The authors wish to thank Andrew Swann for useful discussions and remarks. This research was supported by the Danish Natural Science Research Council, Grant 51-00-0306. The authors are members of the EDGE, Research Training Network HPRN-CT-2000-00101, supported by the European Human Potential Programme. The final part of this paper was done during the visit of S.I. at the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. S.I. thanks the Abdus Salam ICTP for providing support and an excellent research environment. The research of S.I. is partially supported by Contract MM 809/1998 with the Ministry of Science and Education of Bulgaria, Contract 586/2002 with the University of Sofia "St. Kl. Ohridski".

2. GENERAL PROPERTIES OF G_2 -STRUCTURES

We recall some notions of G_2 geometry. Endow $\mathbb{R}^7$ with its standard orientation and inner product. Let $e_1, \dots, e_7$ be an oriented orthonormal basis which we identify with the dual basis via the inner product. Write $e_{i_1 i_2 \dots i_p}$ for the monomial $e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_p}$. We shall omit the $\sum$ -sign understanding summation on any pair of equal indices.

Consider the three-form ω on $\mathbb{R}^7$ given by

$$(2.1) \quad \omega = e_{124} + e_{235} + e_{346} + e_{457} + e_{561} + e_{672} + e_{713}.$$

The subgroup of $GL(7)$ fixing ω is the exceptional Lie group G_2 . It is a compact, connected, simply-connected, simple Lie subgroup of $SO(7)$ of dimension 14 [2].

The Hodge star operator supplies the 4-form $*\omega$ given by

$$(2.2) \quad *\omega = -e_{3567} - e_{4671} - e_{5712} - e_{6123} - e_{7234} - e_{1345} - e_{2456}.$$

We let the expressions

$$\begin{aligned} \omega &= \frac{1}{6} \omega_{ijk} e_{ijk}, \\ *\omega &= \frac{1}{24} \omega_{ijkl} e_{ijkl} \end{aligned}$$

define the symbols ω_{ijk} and ω_{ijkl} . We then obtain the following set of formulae

$$(2.3) \quad \begin{aligned} \omega_{ipq} \omega_{j pq} &= 6 \delta_{ij}, \\ \omega_{ipq} \omega_{jk p q} &= -4 \omega_{ijk}, \\ \omega_{ij p} \omega_{kl p} &= -\omega_{ijkl} + \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}, \\ \omega_{ij p q} \omega_{kl p q} &= -2 \omega_{ijkl} + 4(\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}), \\ \omega_{ij p} \omega_{kl m p} &= \delta_{ik} \omega_{j l m} - \delta_{jk} \omega_{i l m} + \delta_{il} \omega_{j m k} - \delta_{jl} \omega_{i m k} + \delta_{im} \omega_{j k l} - \delta_{jm} \omega_{i k l}. \end{aligned}$$

Definition 2.1. A G_2 -structure on a 7-manifold M is a reduction of the structure group of the tangent bundle to the exceptional group G_2 . Equivalently, there exists a nowhere vanishing differential three-form ω on M and local frames of the cotangent bundle with respect to which ω takes the form (2.1). The three-form ω is called the *fundamental form* of the G_2 -manifold M [1].

We will say that the pair (M, ω) is a G_2 -manifold with G_2 -structure (determined by) ω .

Remark 2.2. Alternatively, a G_2 -structure can be described by the existence of a two-fold vector cross product P on the tangent spaces of M .

The fundamental form of a G_2 -manifold determines a metric through $g_{ij} = \frac{1}{6}\omega_{ikl}\omega_{jkl}$. This is referred to as the metric induced by ω . We write ∇^g for the associated Levi-Civita connection, $||\cdot||^2$ for the tensor norm with respect to g . In addition we will freely identify vectors and co-vectors via the induced metric g .

Let (M, ω) be a G_2 -manifold. The action of G_2 on the tangent space induces an action of G_2 on $\Lambda^k(M)$ splitting the exterior algebra into orthogonal subspaces, where Λ_l^k corresponds to an l -dimensional G_2 -irreducible subspace of Λ^k :

$$\Lambda^1(M) = \Lambda_7^1, \quad \Lambda^2(M) = \Lambda_7^2 \oplus \Lambda_{14}^2, \quad \Lambda^3(M) = \Lambda_1^3 \oplus \Lambda_7^3 \oplus \Lambda_{27}^3,$$

where

$$\begin{aligned} \Lambda_7^2 &= \{\alpha \in \Lambda^2(M) \mid *(\alpha \wedge \omega) = -2\alpha\}, \\ \Lambda_{14}^2 &= \{\alpha \in \Lambda^2(M) \mid *(\alpha \wedge \omega) = \alpha\} \\ \Lambda_1^3 &= \{t\omega \mid t \in \mathbb{R}\}, \\ \Lambda_7^3 &= \{*(\beta \wedge \omega) \mid \beta \in \Lambda^1(M)\}, \\ \Lambda_{27}^3 &= \{\gamma \in \Lambda^3(M) \mid \gamma \wedge \omega = 0, \gamma \wedge *\omega = 0\}. \end{aligned}$$

The Hodge star $*$ gives an isometry between Λ_l^k and Λ_l^{7-k} .

More generally, $V_{(\lambda_1, \lambda_2)}^d$ will denote the G_2 representation of highest weight (λ_1, λ_2) of dimension d . Note that $V_{(0,0)}^1 \cong \Lambda_1^3 \cong \Lambda_1^4$ is the trivial representation, $\Lambda_7^1 \cong V_{(1,0)}^7$ is the standard representation of G_2 on $\mathbb{R}^7$, and the adjoint representation is $\mathfrak{g}_2 \cong V_{(0,1)}^{14} \cong \Lambda_{14}^2$. Also note that $V_{(2,0)}^{27} \cong \Lambda_{27}^3 \cong \Lambda_{27}^4$ is isomorphic to the space of traceless symmetric tensors $S_0^2 V^7$ on $V_{(1,0)}^7$.

3. RICCI TENSORS ON G_2 -MANIFOLD

Let (M, ω) be a G_2 -manifold with fundamental form ω . Let g be the associated Riemannian metric.

$$R_{X,Y} = [\nabla_X^g, \nabla_Y^g] - \nabla_{[X,Y]}^g$$

is then the curvature tensor of the Levi-Civita connection ∇^g of the metric g . The Ricci tensor ρ is defined as usual as the contraction $\rho_{ij} = R_{sij s}$, where $R_{sij s}$ are the components

$$R_{sijk} := g(R(e_s, e_i)e_j, e_k)$$

of the curvature tensor with respect to an orthonormal basis $e_1, \dots, e_7$.

Definition 3.1. On (M, ω) we may define a second symmetric tensor ρ^* by

$$(3.4) \quad \rho_{sm}^* := R_{ijkl}\omega_{ijs}\omega_{klm}.$$

We will call the ρ^* the $\star$ -Ricci tensor of the G_2 -manifold.

The two Ricci tensors have common trace in the following sense. Let $s = \text{tr}_g \rho = \rho_{ii}$ be the scalar curvature and let the trace of ρ^* be denoted by $s^* = \text{tr}_g \rho^* = \rho_{ii}^*$.

Proposition 3.2. *On a G_2 -manifold we have $s^* = -2s$.*

Proof. Apply (2.3) to the definition of s^* and use skew-symmetry of $*\omega$ and the Bianchi identity to conclude that $R_{ijkl}\omega_{ijkl} = 0$. $\square$

Definition 3.3. We shall use the term $\star$ -Einstein for G_2 -manifold (M, ω) when the traceless part of the $\star$ -Ricci tensor vanishes, i.e., when the equation

$$\rho^\star = \frac{s^\star}{7}g$$

holds.

We define associated Ricci three-forms by

$$\begin{aligned}\rho_{ijk}^\star &:= R_{ijlm}\omega_{lmk} + R_{jklm}\omega_{lmi} + R_{kilm}\omega_{lmj}, \\ \rho_{ijk} &:= \rho_{is}\omega_{sjk} + \rho_{js}\omega_{ski} + \rho_{ks}\omega_{sij}.\end{aligned}$$

In terms of the Ricci forms, the Weitzenböck formula for the fundamental form can be written as follows

Proposition 3.4. *On any G_2 -manifold the following formula holds*

$$(3.5) \quad d\delta\omega + \delta d\omega = \nabla^g \star \nabla^g \omega + \rho + \rho^\star.$$

□

4. CLOSED G_2 -STRUCTURES

Let (M^7, ω) be a G_2 -manifold with closed fundamental form. The two-form $\delta\omega$ then takes values in Λ_{14}^2 [3]. As a consequence we get

Proposition 4.1. *The following formulas are valid on a closed G_2 -structure:*

$$(4.6) \quad \delta\omega_{ij}\omega_{ijk} = 0, \quad \delta\omega_{ip}\omega_{pj k} + \delta\omega_{jp}\omega_{pki} + \delta\omega_{kp}\omega_{pij} = 0.$$

□

It is well-known [17] that a G_2 -structure is parallel if and only if it is closed and co-closed, $d\omega = \delta\omega = 0$. The two-form $\delta\omega$ thus may be interpreted as the deviation of ω from a parallel G_2 -structure. We are going to find explicit formulae for the covariant derivatives of the fundamental form of a closed G_2 -structure in terms of $\delta\omega$ and its derivatives.

Definition 4.2. The *canonical connection* $\tilde{\nabla}$ of a closed G_2 -structure may be defined by the equation

$$(4.7) \quad g(\tilde{\nabla}_X Y, Z) = g(\nabla^g_X Y, Z) - \frac{1}{6}\delta\omega(X, e_i)\omega(e_i, Y, Z)$$

for vector fields X, Y, Z .

Using (4.6) it is easy to see that $\tilde{\nabla}$ is a metric G_2 -connection, i.e., it satisfies

$$\tilde{\nabla}\omega = 0, \quad \tilde{\nabla}g = 0.$$

The torsion T of $\tilde{\nabla}$ is determined by

$$g(T(X, Y), Z) = \frac{1}{6}\delta\omega(Z, e_i)\omega(e_i, X, Y).$$

On a compact G_2 -manifold the canonical connection may be characterized as the unique G_2 connection of minimal torsion with respect to the L^2 -norm on M . It may also be described by the fact that the difference $\nabla^g - \tilde{\nabla}$ takes values in Λ_7^2 , the orthogonal complement of $\mathfrak{g}_2 \subset \Lambda^2$ with respect to the metric induced by g .

From the properties of the canonical connection and $\delta\omega$ one derives

Proposition 4.3. *For a closed G_2 -structure the following relations hold:*

$$(4.8) \quad \nabla^g_i \omega_{jkl} = \frac{1}{2} \delta \omega_{ip} \omega_{pjkl},$$

$$(4.9) \quad \nabla^g_i \omega_{jklm} = -\frac{1}{2} (\delta \omega_{ij} \omega_{klm} - \delta \omega_{ik} \omega_{lmj} + \delta \omega_{il} \omega_{mjk} - \delta \omega_{im} \omega_{jkl})$$

and

$$(4.10) \quad \nabla^{g*} \nabla^g \omega_{jkl} = \frac{1}{4} \|\delta \omega\|^2 \omega_{jkl} - \frac{1}{4} (\delta \omega_{ip} \delta \omega_{ij} \omega_{pkl} + \delta \omega_{ip} \delta \omega_{ik} \omega_{plj} + \delta \omega_{ip} \delta \omega_{il} \omega_{pjk}).$$

□

Applying (4.7) and (4.8) we get that the curvature $\tilde{R}$ of the canonical connection $\tilde{\nabla}$ is related to the curvature of the Levi-Civita connection by:

$$(4.11) \quad R_{ijkl} = \tilde{R}_{ijkl} + \frac{1}{6} [\nabla^g_i \delta \omega_{jp} - \nabla^g_j \delta \omega_{ip}] \omega_{pkl} + \frac{1}{9} \delta \omega_{is} \delta \omega_{jp} \omega_{spkl} - \frac{1}{36} [\delta \omega_{ik} \delta \omega_{jl} - \delta \omega_{il} \delta \omega_{jk}]$$

5. CURVATURE OF CLOSED G_2 -STRUCTURES

From here on (M^7, ω) will be a G_2 -manifold with closed G_2 -structure. We have

Proposition 5.1. *The Ricci tensors of a closed G_2 -structure (M, ω) are given by*

$$(5.12) \quad \rho_{lm} = -\frac{1}{4} d\delta \omega_{sjm} \omega_{sjl} + \frac{1}{2} \delta \omega_{lj} \delta \omega_{mj};$$

$$(5.13) \quad \rho_{lm}^* = d\delta \omega_{sjm} \omega_{sjl} + \delta \omega_{lj} \delta \omega_{mj} - \frac{1}{2} \|\delta \omega\|^2 \delta_{ml}.$$

Proof. The Ricci identities for $\omega, * \omega$ together with (4.8) and (4.9) lead to the following useful

Lemma 5.2. *If ω is a closed G_2 -structure on M^7 then*

$$(5.14) \quad \rho_{sr} \omega_{rkl} + \frac{1}{2} R_{skir} \omega_{lir} - \frac{1}{2} R_{slir} \omega_{kir} = -\frac{1}{4} (d\delta \omega_{sjp} + \nabla^g_s \delta \omega_{jp}) \omega_{pjkl} + \frac{1}{2} \delta \omega_{pj} \delta \omega_{sj} \omega_{klp}.$$

$$(5.15) \quad -R_{sijr} \omega_{rkml} - R_{sikr} \omega_{jrml} - R_{silm} \omega_{jkrm} - R_{simr} \omega_{jkrl} = \frac{1}{2} [(\nabla^g_i \delta \omega_{sj} - \nabla^g_s \delta \omega_{ij}) \omega_{klm} - (\nabla^g_i \delta \omega_{sk} - \nabla^g_s \delta \omega_{ik}) \omega_{lmj}] + \frac{1}{2} [(\nabla^g_i \delta \omega_{sl} - \nabla^g_s \delta \omega_{il}) \omega_{mjk} - (\nabla^g_i \delta \omega_{sm} - \nabla^g_s \delta \omega_{im}) \omega_{jkl}] - \frac{1}{4} [(\delta \omega_{ij} \delta \omega_{sp} - \delta \omega_{sj} \delta \omega_{ip}) \omega_{pklm} - (\delta \omega_{ik} \delta \omega_{sp} - \delta \omega_{sk} \delta \omega_{ip}) \omega_{plmj}] - \frac{1}{4} [(\delta \omega_{il} \delta \omega_{sp} - \delta \omega_{sl} \delta \omega_{ip}) \omega_{pmjk} - (\delta \omega_{im} \delta \omega_{sp} - \delta \omega_{sm} \delta \omega_{ip}) \omega_{pjkl}]$$

□

Using (4.7) we get

$$(5.16) \quad \nabla^g_k \delta \omega_{is} \omega_{ism} = \tilde{\nabla}_k \delta \omega_{is} \omega_{ism} + \frac{1}{6} \delta \omega_{kr} \delta \omega_{rq} \omega_{siq} \omega_{ism} = \delta \omega_{kr} \delta \omega_{mr},$$

since $\tilde{\nabla} \delta \omega \in \Lambda_{14}^2$. If we multiply (5.14) by ω_{mkl} and use the Bianchi identity as well as (5.16) we obtain (5.12).

Multiplying (5.15) by ω_{mlj} , and again using the Bianchi identity (alternatively: multiply (4.11) by ω_{klm}), we get

$$(5.17) \quad R_{silm}\omega_{klr} = (\nabla_s^g \delta\omega_{ik} - \nabla_i^g \delta\omega_{sk}) + \frac{1}{4}(\delta\omega_{ij}\delta\omega_{sp} - \delta\omega_{sj}\delta\omega_{ip})\omega_{jpk}.$$

From (5.17) we get that

$$(5.18) \quad \begin{aligned} \rho_{km}^* &= R_{silm}\omega_{klr}\omega_{sim} \\ &= d\delta\omega_{sik}\omega_{sim} - \nabla_k^g \delta\omega_{is}\omega_{sim} + \frac{1}{2}\delta\omega_{ij}\delta\omega_{sp}\omega_{jpk}\omega_{sim}. \end{aligned}$$

The second term is calculated in (5.16). The last term is manipulated using (4.6) and (2.3):

$$(5.19) \quad \begin{aligned} \delta\omega_{ij}\delta\omega_{sp}\omega_{jpk}\omega_{sim} &= (-\delta\omega_{pj}\omega_{jki} - \delta\omega_{kj}\omega_{jip})\delta\omega_{sp}\omega_{sim} \\ &= \delta\omega_{sp}(\delta\omega_{jp}\omega_{kij}\omega_{sim} + \delta\omega_{jk}\omega_{jip}\omega_{sim}) \\ &= \delta\omega_{sp}\delta\omega_{jp}(-\omega_{kjsm} + \delta_{ks}\delta_{jm} - \delta_{km}\delta_{js}) \\ &\quad + \delta\omega_{sp}\delta\omega_{jk}(-\omega_{jpsm} + \delta_{js}\delta_{pm} - \delta_{jm}\delta_{ps}) \\ &= -\|\delta\omega\|^2\delta_{km} + 4\delta\omega_{jm}\delta\omega_{jk}, \end{aligned}$$

again, since $\delta\omega \in \Lambda_{14}^2$. Substituting (5.19) and (5.16) into (5.18) we obtain (5.13). $\square$

The equality (5.16) leads to

$$(5.20) \quad d\delta\omega_{sjm}\omega_{sjm} = 3\|\delta\omega\|^2.$$

Taking the trace in (5.12) and using (5.20), we get the formula for the scalar curvature of a closed G_2 -structure discovered recently by Bryant in [3]

Corollary 5.3. *The scalar curvature of a closed G_2 -structure is non-positive while the $\star$ -scalar curvature is non-negative. These functions are given by*

$$(5.21) \quad s = -\frac{1}{4}\|\delta\omega\|^2, \quad s^* = \frac{1}{2}\|\delta\omega\|^2.$$

In view of (5.21), the trace-free part of the Ricci tensor ρ^0 has the expression

$$(5.22) \quad \rho^0 = \rho + \frac{1}{28}\|\delta\omega\|^2 g.$$

Definition 5.4. The canonical connection gives us a third Ricci tensor which we denote by $\tilde{\rho}$:

$$\tilde{\rho}_{ij} = \tilde{R}_{sij s}.$$

Corollary 5.5. *On a 7-manifold with closed G_2 -structure the Ricci tensor of the canonical connection is related to the Riemannian Ricci tensor through the following formula*

$$\tilde{\rho} = \frac{2}{3}\rho.$$

Proof. Taking the trace of (4.11) we get

$$\rho_{il} = \tilde{\rho}_{il} - \frac{1}{12}d\delta\omega_{pji}\omega_{pjl} + \frac{1}{6}\delta\omega_{is}\delta\omega_{ls}.$$

This equality and (5.12) completes the proof. $\square$

Furthermore, we have

Proposition 5.6. *The Ricci three-forms of a closed G_2 -structure are given by*

$$(5.23) \quad \rho_{jlk} = -\frac{1}{4} \|\delta\omega\|^2 \omega_{jlk} - d\delta\omega_{jlk} - \frac{1}{2} (\delta\omega_{li} \delta\omega_{pi} \omega_{pjk} + \delta\omega_{ki} \delta\omega_{pi} \omega_{plj} + \delta\omega_{ji} \delta\omega_{pi} \omega_{pkl})$$

$$(5.24) \quad \rho_{sik}^* = 2d\delta\omega_{sik} + \frac{1}{4} (\delta\omega_{ij} \delta\omega_{pj} \omega_{psk} + \delta\omega_{kj} \delta\omega_{pj} \omega_{pis} + \delta\omega_{sj} \delta\omega_{pj} \omega_{pki}).$$

Proof. Substitute (4.10) and (5.24) into the Weitzenböck formula (3.5) to get (5.23). The cyclic sum in the equality (5.17) gives (5.24). $\square$

Theorem 5.7. *Let (M^7, ω) be a G_2 -manifold with closed fundamental form. If (M, ω) is Einstein and $\star$ -Einstein then M is parallel.*

Proof. Let (M, ω) be a G_2 manifold with closed G_2 structure ω . Suppose that both the Einstein and $\star$ -Einstein equations are satisfied. Proposition 5.1 in this case yields

$$(5.25) \quad \delta\omega_{ij} \delta\omega_{pj} = \frac{1}{7} \|\delta\omega\|^2 \delta_{ip}, \quad d\delta\omega_{ijk} \omega_{ijm} = \frac{3}{7} \|\delta\omega\|^2 \delta_{km}.$$

Taking into account (5.25) and the equalities (4.10), (5.23), we obtain

$$(5.26) \quad \begin{aligned} \rho_{ijk}^* &= 2d\delta\omega_{ijk} - \frac{3}{28} \|\delta\omega\|^2 \omega_{ijk}, \\ \nabla^{g*} \nabla^g \omega_{ijk} &= \frac{1}{7} \|\delta\omega\|^2 \omega_{ijk}, \\ \rho_{ijk} &= -\frac{3}{28} \|\delta\omega\|^2 \omega_{ijk}. \end{aligned}$$

In view of (5.26), the Weitzenböck formula (3.5) gives

$$(5.27) \quad d\delta\omega = \frac{1}{14} \|\delta\omega\|^2 \omega.$$

Bryant shows in [3] that on an Einstein manifold with closed G_2 -structure the Λ_{27}^3 -part of $d\delta\omega$ is given by the Λ_{27}^3 -part of $*(\delta\omega \wedge \delta\omega)$. Comparing with (5.27), we see that the Λ_{27}^3 -part of $*(\delta\omega \wedge \delta\omega)$ vanishes. We need the following algebraic

Lemma 5.8. *Let α be a two-form in Λ_{14}^2 . Then the Λ_{27}^3 -part of $*(\alpha \wedge \alpha)$ vanishes if and only if $\alpha = 0$.*

Proof. First note that if α is a two-form in Λ_{14}^2 then $\alpha \otimes \alpha \in S^2 \Lambda_{14}^2 \subset S^2(\Lambda^2)$. The space of symmetric tensors on Λ_{14}^2 decomposes as follows

$$S^2 \Lambda_{14}^2 = V_{(0,2)}^{77} + V_{(0,2)}^{27} + V_{(0,0)}^1.$$

Recall that the map $S^2 \Lambda^2 \rightarrow \Lambda^4$ given by $\beta \vee \gamma \mapsto \beta \wedge \gamma$ is surjective and equivariant. By Schurs Lemma we may conclude that $\alpha \wedge \alpha \in \Lambda_{27}^4 \oplus \Lambda_1^4$ if $\alpha \in \Lambda_{14}^2$.

Now suppose $(\alpha \wedge \alpha)_{\Lambda_{27}^4} = 0$. We may then conclude that $\alpha \wedge \alpha = c\omega$ for some constant c . However, as α is a two-form on an odd-dimensional space it is degenerate. Let $X \in \mathbb{R}^7$ be a non-zero vector such that $X \lrcorner \alpha = 0$. Then

$$cX \lrcorner \omega = X \lrcorner (\alpha \wedge \alpha) = 2(X \lrcorner \alpha) \wedge \alpha = 0.$$

But the left hand side vanishes only if $c = 0$.

This shows that $\alpha \otimes \alpha \in V_{(0,2)}^{77}$ whenever $\alpha \in \Lambda_{14}^2$ satisfies $(\alpha \wedge \alpha)_{\Lambda_{27}^4} = 0$. But $V_{(0,2)}^{77}$ is precisely the space in which the Ricci-flat Riemannian curvature of parallel G_2 -manifolds takes values, whence $\|\alpha\|^2 = 0$. $\square$

Now, Lemma 5.8 implies $\delta\omega = 0$, whence $\nabla^g \omega = 0$. $\square$

Remark 5.9. R.L.Bryant observe [4] that the following identity holds

$$(5.28) \quad \|\alpha \wedge \alpha\|^2 = 6 \|\alpha\|^4, \quad \alpha \in \Lambda_{14}^2.$$

Clearly (5.28) implies the lemma. Note that the constants have been changed to fit our conventions.

6. AN INTEGRAL FORMULA ON CLOSED G_2 MANIFOLD

Our main technical tool to handle the closed G_2 -structure on a compact manifold is the next

Proposition 6.1. *Let (M, ω, g) be a compact G_2 -manifold with closed fundamental form. Then the following integral formula holds:*

$$(6.29) \quad \int_M \left(\frac{1}{24} \|\delta\omega\|^4 + \frac{28}{9} \|\rho^0\|^2 - \frac{7}{18} \rho_{pl}^0 \delta\omega_{bl} \delta\omega_{bp} - \frac{7}{9} (\delta R)_{bkl} \omega_{jkl} \delta\omega_{bj} \right) dV = 0$$

Proof. The first Pontrjagin form $p_1(\nabla)$ of a connection ∇ may be defined by

$$p_1(\nabla) := \frac{1}{16\pi^2} R_{ijab} R_{klab} e_i \wedge e_j \wedge e_k \wedge e_l.$$

The first Pontrjagin class of TM which is the de Rham cohomology class whose representative element is the first Pontrjagin form of a some affine connection on M is independent of the connection on M . This implies that $(p_1(\nabla^g) - p_1(\tilde{\nabla}))$ is an exact 4-form. Since the fundamental form ω is closed, the wedge product $(p_1(\nabla^g) - p_1(\tilde{\nabla})) \wedge \omega$ is exact. From Stoke's theorem we obtain

$$(6.30) \quad \int_M (p_1(\nabla^g) - p_1(\tilde{\nabla})) \wedge \omega = 0.$$

However, we may also express the integrand in terms of the curvatures of ∇^g and $\tilde{\nabla}$ using that

$$16\pi^2 p_1(\nabla^g) \wedge \omega = R_{ijab} R_{klab} \omega_{ijkl} = R_{abij} R_{klab} \omega_{ijkl},$$

and

$$16\pi^2 p_1(\tilde{\nabla}) \wedge \omega = \tilde{R}_{ijab} \tilde{R}_{klab} \omega_{ijkl}.$$

From this point on the proof is essentially a brute force calculation reducing the difference of these two expressions to the form (6.29).

Since $\tilde{\nabla}$ is a G_2 -connection we get

$$(6.31) \quad \tilde{R}_{abij} \omega_{ijkl} = 2\tilde{R}_{abkl}.$$

Using (4.11), (2.3) and (6.31) we calculate

$$(6.32) \quad 16\pi^2 p_1(\nabla^g) \wedge \omega = 2R_{klab} \tilde{R}_{abkl} - \frac{2}{3} (d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab}) \omega_{pkl} R_{klab} \\ - \frac{5}{18} \delta\omega_{ai} \delta\omega_{bj} \omega_{ijkl} R_{klab} + \frac{8}{9} \delta\omega_{ak} \delta\omega_{bl} R_{klab}.$$

Applying (4.11) to (6.32) and using (2.3) we obtain after some calculations that

$$(6.33) \quad 16\pi^2 p_1(\nabla^g) \wedge \omega = \\ 2\tilde{R}_{klab} \tilde{R}_{abkl} + (d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab}) \omega_{pkl} \left(-\frac{2}{3} R_{klab} + \frac{1}{3} \tilde{R}_{klab} \right) \\ + \delta\omega_{ai} \delta\omega_{bj} \omega_{ijkl} \left(-\frac{5}{18} R_{klab} + \frac{2}{9} \tilde{R}_{klab} \right) + \delta\omega_{ak} \delta\omega_{bl} \left(\frac{8}{9} R_{klab} - \frac{1}{9} \tilde{R}_{klab} \right).$$

To calculate the $p_1(\tilde{\nabla})$ term of the integrand we first observe that (4.11) implies

$$(6.34) \quad \tilde{R}_{ijkl} = \tilde{R}_{klij} + \frac{1}{6}(d\delta\omega_{klp} - \nabla_p^g \delta\omega_{kl})\omega_{pij} \\ - \frac{1}{6}(d\delta\omega_{ijp} - \nabla_p^g \delta\omega_{ij})\omega_{pkl} + \frac{1}{9}\delta\omega_{ks}\delta\omega_{lp}\omega_{spij} - \frac{1}{9}\delta\omega_{is}\delta\omega_{jp}\omega_{spkl}.$$

Taking into account (6.31), (6.34) and (2.3), we calculate

$$(6.35) \quad 16\pi^2 p_1(\tilde{\nabla}) \wedge \omega = 2\tilde{R}_{klab}\tilde{R}_{abkl} - \frac{2}{3}(d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab})\omega_{pkl}\tilde{R}_{klab} \\ - \frac{4}{9}\delta\omega_{ai}\delta\omega_{bj}\omega_{ijkl}\tilde{R}_{klab} + \frac{8}{9}\delta\omega_{ak}\delta\omega_{bl}\tilde{R}_{klab}.$$

Subtracting (6.35) from (6.33) we get

$$(6.36) \quad 16\pi^2(p_1(\nabla^g) - p_1(\tilde{\nabla})) \wedge \omega = A + B + C,$$

where A, B, C are given by

$$(6.37) \quad A = (d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab})\omega_{pkl}\left(\frac{1}{3}R_{klab} - (R_{klab} - \tilde{R}_{klab})\right)$$

$$(6.38) \quad B = \delta\omega_{ai}\delta\omega_{bj}\omega_{ijkl}\left(\frac{7}{18}R_{klab} - \frac{2}{3}(R_{klab} - \tilde{R}_{klab})\right)$$

$$(6.39) \quad C = \delta\omega_{ak}\delta\omega_{bl}\left(-\frac{1}{9}R_{klab} + (R_{klab} - \tilde{R}_{klab})\right).$$

We shall calculate each term in (6.36).

First observe that (5.28) implies the useful identity

$$(6.40) \quad 4\delta\omega_{ai}\delta\omega_{bi}\delta\omega_{aj}\delta\omega_{bj} = \|\delta\omega\|^4.$$

Taking into account (4.11), (5.17), (5.12), (5.16), (5.19), (5.20) and (6.40) we obtain after some calculation that

$$(6.41) \quad A = -\frac{19}{24.7}\|\delta\omega\|^4 + \frac{1}{3}\|d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab}\|^2 - \frac{8}{3}\|\rho^0\|^2 + \frac{4}{3}\rho_{sp}^0\delta\omega_{st}\delta\omega_{pt} \\ - \frac{1}{9}(d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab})\omega_{ijp}\delta\omega_{ai}\delta\omega_{bj} - \frac{1}{9}(d\delta\omega_{abp} - \nabla_p^g \delta\omega_{ab})\omega_{pkl}\omega_{smab}\delta\omega_{ks}\delta\omega_{lm}.$$

Let $X_s = R_{akbl}\omega_{sak}\delta\omega_{bl}$, $Y_s = \delta\omega_{bl}\delta\omega_{li}\delta\omega_{bp}\omega_{ips}$. Using the first and second Bianchi identity as well as (5.17), we get

$$(6.42) \quad R_{klab}\delta\omega_{ak}\delta\omega_{bl} = \frac{1}{2}R_{akbl}\delta\omega_{ak}\delta\omega_{bl} \\ = -\frac{1}{2}R_{akbl}\nabla_s^g \omega_{sak}\delta\omega_{bl} \\ = \frac{1}{2}\delta X + \frac{1}{2}R_{akbl}\omega_{sak}\nabla_s^g \delta\omega_{bl} \\ = \frac{1}{2}\delta X + \frac{1}{6}\|d\delta\omega\|^2 - \frac{1}{2}\|\nabla^g \delta\omega\|^2 + \frac{1}{4}\nabla_s^g \delta\omega_{bl}\delta\omega_{li}\delta\omega_{bp}\omega_{ips} \\ = \frac{1}{2}\delta X - \frac{1}{4}\delta Y + \frac{1}{6}\|d\delta\omega\|^2 - \frac{1}{2}\|\nabla^g \delta\omega\|^2 \\ - \frac{1}{8}\|\delta\omega\|^4 + \frac{1}{4}d\delta\omega_{sil}\delta\omega_{bl}\delta\omega_{bp}\omega_{ips}.$$

Applying (5.12) to (6.42), we obtain

$$(6.43) \quad R_{klab}\delta\omega_{ak}\delta\omega_{bl} = \frac{1}{2}\delta X - \frac{1}{4}\delta Y + \frac{1}{6}\|d\delta\omega\|^2 - \frac{1}{2}\|\nabla^g \delta\omega\|^2 + \frac{1}{28}\|\delta\omega\|^4 - \rho_{sp}^0\delta\omega_{st}\delta\omega_{pt}.$$

Remark 6.2. Notice that (6.43) is the Weitzenböck formula

$$\delta d\delta\omega = \nabla^{g*}\nabla^g\delta\omega - \frac{1}{14}\|\delta\omega\|^2\delta\omega + \rho^0(\delta\omega, \cdot) + \rho^0(\cdot, \delta\omega) + R(\delta\omega)$$

for the 2-form $\delta\omega$ on a closed G_2 -structure.

Using (4.11), (5.28) and (6.43) we get that

$$(6.44) \quad C = -\frac{1}{36}\delta(2X - Y) - \frac{25}{63.16}\|\delta\omega\|^4 - \frac{1}{54}\|d\delta\omega\|^2 + \frac{1}{18}\|\nabla^g\delta\omega\|^2 \\ + \frac{1}{9}\rho_{sp}^0\delta\omega_{st}\delta\omega_{pt} + \frac{1}{6}(d\delta\omega_{kls} - \nabla^g_s\delta\omega_{kl})\omega_{sab}\delta\omega_{ak}\delta\omega_{bl}.$$

Now we calculate B . Denote $Z_a = R_{klab}\omega_{jkl}\delta\omega_{bj}$. Using (4.8) and (5.17) we have the following sequence of equalities

$$(6.45) \quad \frac{1}{2}R_{klab}\delta\omega_{ai}\omega_{ijkl}\delta\omega_{bj} = R_{klab}\nabla^g_a\omega_{jkl}\delta\omega_{bj} \\ = -\delta Z + (\delta R)_{bkl}\omega_{jkl}\delta\omega_{bj} - R_{klab}\omega_{jkl}\nabla^g_a\delta\omega_{bj} \\ = -\delta Z + (\delta R)_{bkl}\omega_{jkl}\delta\omega_{bj} - \frac{1}{6}\|d\delta\omega\|^2 - \frac{1}{2}\|\nabla^g\delta\omega\|^2 \\ + \frac{1}{4}(d\delta\omega_{abj} - \nabla^g_j\delta\omega_{ab})\omega_{slj}\delta\omega_{as}\delta\omega_{bl}$$

Applying (4.11), we get

$$(6.46) \quad -\frac{2}{3}(R_{klab} - \tilde{R}_{klab})\delta\omega_{ai}\delta\omega_{bj}\omega_{ijkl} = \\ -\frac{1}{9}(d\delta\omega_{kls} - \nabla^g_s\delta\omega_{kl})\omega_{sab}\omega_{ijkl}\delta\omega_{ai}\delta\omega_{bj} \\ -\frac{2}{27}\delta\omega_{ks}\delta\omega_{lm}\delta\omega_{ai}\delta\omega_{bj}\omega_{smab}\omega_{ijkl}.$$

Denote $V_k = \omega_{mab}\delta\omega_{lm}\delta\omega_{bj}\delta\omega_{ai}\omega_{ijkl}$. Using (4.8), we obtain

$$(6.47) \quad \frac{1}{2}\delta\omega_{ks}\omega_{smab}\delta\omega_{lm}\delta\omega_{ai}\delta\omega_{bj}\omega_{ijkl} = \nabla^g_k\omega_{mab}\delta\omega_{lm}\delta\omega_{ai}\delta\omega_{bj}\omega_{ijkl} \\ = -\delta V - \frac{3}{2}(d\delta\omega_{klm} - \nabla^g_m\delta\omega_{kl})\omega_{mab}\omega_{ijkl}\delta\omega_{ai}\delta\omega_{bj}.$$

Get together (6.45), (6.46) and (6.47) we obtain

$$(6.48) \quad B = -\frac{1}{27}\delta(21Z - 4V) + \frac{7}{9}(\delta R)_{bkl}\omega_{jkl}\delta\omega_{bj} - \frac{7}{54}\|d\delta\omega\|^2 - \frac{7}{18}\|\nabla^g\delta\omega\|^2 \\ + \frac{1}{9}(d\delta\omega_{klm} - \nabla^g_m\delta\omega_{kl})\omega_{mab}\omega_{ijkl}\delta\omega_{ai}\delta\omega_{bj} \\ + \frac{7}{36}(d\delta\omega_{abj} - \nabla^g_j\delta\omega_{ab})\omega_{slj}\delta\omega_{as}\delta\omega_{bl}.$$

Collecting terms from (6.41), (6.44) and (6.48) we obtain the following expression

$$(6.49) \quad 16\pi^2(p_1(\nabla^g) - p_1(\tilde{\nabla})) \wedge \omega = \delta\left(-\frac{1}{18}X + \frac{1}{36}Y - \frac{7}{9}Z + \frac{4}{27}V\right) \\ - \frac{139}{18.56}\|\delta\omega\|^4 - \frac{1}{27}\|d\delta\omega\|^2 - \frac{8}{3}\|\rho^0\|^2 \\ + \frac{7}{9}(\delta R)_{bkl}\omega_{jkl}\delta\omega_{bj} + \frac{1}{4}(d\delta\omega_{kls} - \nabla^g_s\delta\omega_{kl})\omega_{sab}\delta\omega_{ak}\delta\omega_{bl} + \frac{13}{9}\rho_{pt}^0\delta\omega_{bl}\delta\omega_{bp}.$$

We express the last two terms in a more tractable form. Applying (4.6) we have

$$\begin{aligned} d\delta\omega_{abp}\delta\omega_{ai}\delta\omega_{bj}\omega_{pij} &= -d\delta\omega_{abp}\delta\omega_{ai}(\delta\omega_{pj}\omega_{ibj} + \delta\omega_{ij}\omega_{bpij}) \\ &= -d\delta\omega_{apb}\delta\omega_{ai}\delta\omega_{bj}\omega_{ipj} - d\delta\omega_{abp}\delta\omega_{ai}\delta\omega_{ij}\omega_{bpij}, \end{aligned}$$

whence

$$(6.50) \quad d\delta\omega_{abp}\delta\omega_{ai}\delta\omega_{bj}\omega_{pij} = \frac{1}{2}d\delta\omega_{abp}\omega_{bpij}\delta\omega_{ai}\delta\omega_{ji} = -2\rho_{aj}\delta\omega_{ai}\delta\omega_{ji} + \frac{1}{4}\|\delta\omega\|^4,$$

where we used (6.40) and (5.12).

The next equalities are a consequence of (5.12), (5.28) and (5.16)

$$\begin{aligned} (6.51) \quad \nabla^g_s \delta\omega_{kl}\omega_{sab}\delta\omega_{ak}\delta\omega_{bl} &= \delta Y + \frac{1}{4}\|\delta\omega\|^4 - (d\delta\omega_{sak} - \nabla^g_a \delta\omega_{ks})\omega_{sab}\delta\omega_{kl}\delta\omega_{bl} \\ &= \delta Y + 4\rho_{kb}\delta\omega_{kl}\delta\omega_{bl}. \end{aligned}$$

Substitute (6.50), (6.51) into (6.49), use (5.12) and integrate over the compact M to get

$$(6.52) \quad \int_M \left(-\frac{11}{18 \cdot 28} \|\delta\omega\|^4 - \frac{1}{27} \|d\delta\omega\|^2 - \frac{8}{3} \|\rho^0\|^2 - \frac{1}{18} \rho_{pl}^0 \delta\omega_{bl}\delta\omega_{bp} + \frac{7}{9} (\delta R)_{bkl}\omega_{jkl}\delta\omega_{bj} \right) dV = 0.$$

We calculate from (5.23) that

$$(6.53) \quad \|d\delta\omega\|^2 = \frac{15}{28} \|\delta\omega\|^4 + 12 \|\rho^0\|^2 - 12\rho_{pl}^0 \delta\omega_{bl}\delta\omega_{bp}.$$

Substitute (6.53) into (6.52) to obtain (6.29). $\square$

Corollary 6.3. (Integral Weitzenböck formula.) *On a compact G_2 manifold with closed fundamental form we have*

$$\int_M R_{akbl}\delta\omega_{ak}\delta\omega_{bl} dV = \int_M \left(\frac{1}{3} \|d\delta\omega\|^2 - \|\nabla^g \delta\omega\|^2 - 2\rho_{sp}\delta\omega_{st}\delta\omega_{pt} \right) dV.$$

Proof. The first Bianchi identity implies $R_{akbl}\delta\omega_{ak}\delta\omega_{bl} = 2R_{klab}\delta\omega_{ak}\delta\omega_{bl}$. Apply (5.22) to (6.43) multiply by two and integrate the obtained equality over the compact manifold to get the result. $\square$

7. PROOF OF THE MAIN THEOREM

We consider the co-differential of the Weyl tensor, δW as an element of $T^*M \otimes \Lambda^2(T^*M)$. According to the splitting of the space of two forms, δW splits as follows

$$\delta W = \delta W_{14}^2 \oplus \delta W_7^2,$$

where δW_{14}^2 is a section of $T^*M \otimes \Lambda_{14}^2(T^*M) \cong \mathbb{R}^7 \otimes g_2$ while δW_7^2 is a section of $T^*M \otimes \Lambda_7^2(T^*M)$.

The Main Theorem is a consequence of the following general

Theorem 7.1. *Let (M, ω, g) be a compact G_2 -manifold with closed fundamental form ω . Suppose that the Λ_7^2 -part of the co-differential of the Weyl tensor vanishes, $\delta W = \delta W_{14}^2$. Then (M, ω, g) is a Joyce space*

Proof. On a 7-dimensional Riemannian manifold the Weyl tensor is expressed in terms of the normalized Ricci tensor $h = -\frac{1}{5}(\rho - \frac{s}{12}g)$ as follows

$$W_{ijkl} = R_{ijkl}^g - h_{ik}g_{jl} + h_{jk}g_{il} - h_{jl}g_{ik} + h_{il}g_{jk}.$$

The second Bianchi identity implies

$$(7.54) \quad \nabla^g_i \rho_{jk} - \nabla^g_j \rho_{ik} = (\delta R)_{kij}, \quad ds_i = 2\nabla^g_j \rho_{ji}, \quad (\delta W)_{kij} = -4(\nabla^g_i h_{jk} - \nabla^g_j h_{ik}).$$

The condition of the theorem reads

$$0 = (\delta W)_{kij} \omega_{ijl} = \frac{4}{5}(\nabla^g_i \rho_{jk} - \nabla^g_j \rho_{ik}) \omega_{ijl} - \frac{2}{15} ds_i \omega_{ikl}.$$

Consequently,

$$(7.55) \quad \frac{4}{5}(\delta R)_{kij} \omega_{ijl} \delta \omega_{kl} = (\nabla^g_i \rho_{jk} - \nabla^g_j \rho_{ik}) \omega_{ijl} \delta \omega_{kl} = 0,$$

since $\delta \omega \in \Lambda_{14}^2$.

To apply effectively our integral formula (6.29) we have to evaluate one more term.

Denote $K_i = \omega_{ijl} \rho_{jk} \delta \omega_{kl}$. The equation (7.55) together with (5.12) leads to the equality

$$\delta K = -\frac{1}{2} \rho_{jk} \delta \omega_{jl} \delta \omega_{kl} - 2\|\rho\|^2$$

which implies, by an integration over the compact M , that

$$(7.56) \quad \int_M \rho_{jk} \delta \omega_{jl} \delta \omega_{kl} dV = -4 \int_M \|\rho\|^2 dV.$$

The equalities (7.55), (7.56), (5.22) and (6.29) yield

$$\int_M \left(\frac{1}{24} \|\delta \omega\|^4 + \frac{14}{3} \|\rho^0\|^2 \right) dV = 0.$$

Hence, Theorem 7.1 follows. □

Clearly, our Main Theorem follows from Theorem 7.1. □

Corollary 5.5 and the main theorem lead to

Theorem 7.2. *Any compact 7-manifold with closed G_2 -structure which is Einstein with respect to the canonical connection is a Joyce space.*

REFERENCES

- [1] E. Bonan, *Sur les variétés riemanniennes à groupe d'holonomie G_2 ou $Spin(7)$* , C. R. Acad. Sci. Paris **262** (1966), 127-129. 1, 3
- [2] R. Bryant, *Metrics with exceptional holonomy*, Ann. Math. **126** (1987), 525-576. 1, 3
- [3] ———, *Some remarks on G_2 -structures*, math.DG/0305124, Version 1. 2, 5, 7, 8
- [4] ———, *Some remarks on G_2 -structures*, math.DG/0305124, Version 2. 2, 9
- [5] R. Bryant, S. Salamon, *On the construction of some complete metrics with exceptional holonomy*, Duke Math. J. **58** (1989), 829-850. 2
- [6] F. Cabrera, M. Monar, A. Swann, *Classification of G_2 -structures*, J. London Math. Soc. **53** (1996), 407-416. 2
- [7] S. Chiossi, S. Salamon, *The intrinsic torsion of $SU(3)$ and G_2 -structures*, Differential Geometry, Valencia 2001, World Sci. Publishing, 2002, pp. 115-133, math.DG/0202282. 1
- [8] R. Cleyton, S. Ivanov, *On the geometry of closed G_2 -structures*, math.DG/0306362, Version 1, IMADA preprint PP-2003-11. 2
- [9] M. Cvetič, G.W. Gibbons, H. Lu, C.N. Pope, *Almost Special Holonomy in Type IIA/M Theory*, Nucl. Phys. **B638** (2002), 186-206, hep-th/0203060. 2
- [10] M. Fernandez, *An Example of compact calibrated manifold associated with the exceptional Lie group G_2* , J. Diff. Geom. **26** (1987), 367-370. 2

- [11] M. Fernandez, A. Gray, *Riemannian manifolds with structure group G_2* , Ann. Mat. Pura Appl. (4) **32** (1982), 19-45. 1
- [12] Th. Friedrich, S. Ivanov *Parallel spinors and connections with skew-symmetric torsion in string theory*, Asian J. Math. **6** (2002), 303-336, math.DG/0102142 . 2
- [13] Th. Friedrich, S. Ivanov, *Killing spinor equations in dimension 7 and geometry of integrable G_2 manifolds*, J. Geom. Phys. **48** (2003), 1-11, math.DG/0111131. 2
- [14] J. Gauntlett, N. Kim, D. Martelli, D. Waldram, *Fivebranes wrapped on SLAG three-cycles and related geometry*, JHEP 0111 (2001) 018, hep-th/0110034. 2
- [15] J. Gauntlett, D. Martelli, D. Waldram, *Superstrings with Intrinsic torsion*, Phys. Rev. **D69** (2004) 086002, hep-th/0302158. 2
- [16] G.W. Gibbons, D.N. Page, C.N. Pope, *Einstein metrics on S^3 , $\mathbb{R}^3$, and $\mathbb{R}^4$ bundles*, Commun. Math. Phys. **127** (1990), 529-553. 2
- [17] A. Gray, *Vector cross product on manifolds*, Trans. Am. Math. Soc. **141** (1969), 463-504, Correction **148** (1970), 625. 1, 5
- [18] P. Ivanov, S. Ivanov, *$SU(3)$ -instantons and G_2 , $Spin(7)$ -heterotic string solitons*, Commun. Math. Phys., in press, math.DG/0312094. 2
- [19] D. Joyce, *Compact Riemannian 7-manifolds with holonomy G_2 . I*, J. Diff. Geom. **43** (1996), 291-328. 2
- [20] ———, *Compact Riemannian 7-manifolds with holonomy G_2 . II*, J. Diff. Geom., **43** (1996), 329-375. 2
- [21] ———, *Compact Riemannian manifolds with special holonomy*, Oxford University Press, 2000. 2
- [22] A. Kovalev, *Twisted connected sums and special Riemannian holonomy*, J. Reine Angew. math. **565** (2003), 125-160, math.DG/0012189. 2
- [23] B. Lawson, M.-L. Michelsohn, *Spin Geometry*, Princeton University Press, 1989. 1
- [24] N. C. Leung, *TQFT for Calabi-Yau three folds and G_2 manifolds*, Adv. Theor. Math. Phys. **6** (2002) 575-591, math.DG/0208124. 2
- [25] S. Salamon, *Riemannian geometry and holonomy groups*, Pitman Res. Notes Math. Ser., 201 (1989). 1

(Cleyton) DEPARTMENT OF MATHEMATICS, SURGE 222, UNIVERSITY OF CALIFORNIA RIVERSIDE, CA 92521 - USA

E-mail address: cleyton@math.ucr.edu

(Ivanov) UNIVERSITY OF SOFIA "ST. KL. OHRIDSKI", FACULTY OF MATHEMATICS AND INFORMATICS,, BLVD. JAMES BOURCHIER 5,, 1164 SOFIA, BULGARIA

E-mail address: ivanovsp@fmi.uni-sofia.bg